



Venom proteomic analysis of medically important Nigerian viper *Echis ocellatus* and *Bitis arietans* snake species

Emeka John Dingwoke^{a,f}, Fatima Amin Adamude^{b,f}, Gadija Mohamed^c, Ashwil Klein^d, Aliyu Salihu^a, Mujitaba Suleiman Abubakar^{e,f}, Abdullahi Balarabe Sallau^{a,f,*}

^a Department of Biochemistry, Faculty of Life Sciences, Ahmadu Bello University, Zaria, Kaduna State, Nigeria

^b Department of Biochemistry, Faculty of Sciences, Federal University, Lafia, Nasarawa State, Nigeria

^c Agri-Food Systems and Omics, Post-Harvest and Agro-Processing Technologies, Agricultural Research Council, Infrutec-Nietvoorbij, Stellenbosch, 7599, South Africa

^d Department of Biotechnology, Faculty of Natural Sciences, Proteomics Research Unit, University of Western Cape, South Africa

^e Department of Pharmacognosy and Drug Development, Faculty of Pharmaceutical Sciences, Ahmadu Bello University, Zaria, Kaduna State, Nigeria

^f Venom, Antivenom and Natural Toxins Research Centre, Ahmadu Bello University, Zaria, Kaduna State, Nigeria

ARTICLE INFO

Keywords:

Snakebite
Neglected tropical disease
Venom-based antivenom
Viperidae
Bitis arietans
Echis ocellatus
Snake venom
Proteomic
Mass spectrometry

ABSTRACT

Snakebite envenoming remains a neglected tropical disease which poses severe health hazard, especially for the rural inhabitants in Africa. In Nigeria, vipers are responsible for the highest number of deaths. Hydrophilic interaction liquid chromatography coupled with LC-MS/MS was used to analyze the crude venoms of *Echis ocellatus* (Carpet viper) and *Bitis arietans* (Puff adder) in order to understand their venom proteomic identities. Results obtained revealed that gel-free proteomic analysis of the crude venoms led to the identification of 85 and 79 proteins, respectively. Seventy-eight (78) proteins were common between the two snake species with a 91.8% similarity score. The identified proteins belong to 18 protein families in *E. ocellatus* and 14 protein families in *B. arietans*. Serine proteases (22.31%) and metalloproteinases (21.06%) were the dominant proteins in the venom of *B. arietans*; while metalloproteinases (34.84%), phospholipase A₂s (21.19%) and serine proteases (15.50%) represent the major toxins in the *E. ocellatus* venom. Other protein families such as three-finger toxins and cysteine-rich venom proteins were detected in low proportions. This study provides an insight into the venom proteomic analysis of the two Nigerian viper species, which could be useful in identifying the toxin families to be neutralized in case of envenomation.

1. Introduction

Venomous snakes pose severe health problems especially for the rural dwellers in the tropical regions of developing countries [1–3]. Despite the remarkable efforts for effective management of envenomings, the World Health Organization (WHO) incorporated snakebite in the list of neglected tropical diseases [4]. Thus, further and better efforts are required to address the life-threatening situation posed by this disease. Snakebites are a common health hazard in the Savanna region of West Africa [2–4]. In Nigeria, the *Viperidae*, notably *Bitis arietans* (Puff adder) and *Echis ocellatus* (Carpet viper) are associated with the highest incidence of morbidity and mortality [1,5–7] and are regarded as the most dangerous venomous snake species. The actual fatality index resulting from envenomings by these vipers remains unknown due to

inaccurate epidemiological data [1,2,8], as most of the victims do not have access to health care facilities and therefore resort to ethno-medicinal means for treatments. Reports from community-based surveys suggest a higher envenomation incidence than the hospital-based estimates [2,8,9]. The WHO reported about 5.4 million incidences of snakebites annually, with 2.7 million cases of envenomings, 138,000 deaths and 400,000 cases of disabilities [10]. In Nigeria, the incidence of snakebites leads to 10,000 deaths annually where envenomation by *E. ocellatus* was responsible for 66% of total cases [7,11]. Snakebite inflicts severe damage to different organs of the body, and envenomings by the *Viperidae* are characterized by life-threatening symptoms, primarily due to local tissue damage and bleeding [1].

The evolution of venomomics and antivenomics using the ‘-omics’ technologies has paved a way to obtain valuable insights into snake

* Corresponding author. Department of Biochemistry/Venom, Antivenom and Natural Toxins Research Centre, Ahmadu Bello University, Zaria, Kaduna State, Nigeria.

E-mail address: bsallau@yahoo.com (A.B. Sallau).

<https://doi.org/10.1016/j.bbrep.2021.101164>

Received 8 September 2021; Received in revised form 13 October 2021; Accepted 25 October 2021

2405-5808/© 2021 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. One-dimensional SDS-PAGE profile of *B. arietans* and *E. ocellatus* venoms.

venom compositions, revealing the protein constituents and different types of toxic components [12,13]. In addition, the ‘-omics’ approach has led to the rapid examination of the immune reactivity of antivenoms against the toxins found in snake venoms [2,14]. Due to the complex nature of the protein mixture in snake venoms, envenomation is characterized by varied pathological manifestations [2]. It is reported that geographical inhabitation of snake [15–18], diet type [19] and age [20] contribute to the variability of the venom composition. Hence, analyzing venom proteomes to gain an insight into snake venom compositions can help in understanding their biological and pathological effects [21]. It will also help in making appropriate choice of antivenom to use in treatment. The venom gland transcriptomic of Nigerian *B. arietans* and *E. ocellatus* was reported previously [22,23]. In the current study, the venom proteomes of *B. arietans* and *E. ocellatus* captured from different regions of Nigeria were analyzed with the view to understanding their toxin profile and protein identities.

2. Materials and method

2.1. *Bitis arietans* and *Echis ocellatus* venom samples

For this study, 42 male snakes (36 *E. ocellatus* and 6 *B. arietans*) were used. All protocols for snake handling were observed and carried out in accordance with the Ahmadu Bello University Committee on Animal Use and Care and in compliance with the revised ARRIVE guidelines 2.0. Adult snakes *B. arietans* and *E. ocellatus* were captured from different regions of the Southern and Northern parts of Nigeria. This sampling strategy justifies a generalized inference about the protein profile of

these Nigerian vipers. The snakes were maintained at the serpentarium of the Department of Veterinary Pharmacology and Toxicology of Ahmadu Bello University. After 3 days, their venoms were manually extracted as described by Hill and Mackessy [24]. Venoms from snakes of the same species were combined. They were frozen at -80°C , lyophilized using a freeze-dryer, and stored at -20°C .

2.2. Chemicals and reagents

Magnetic bead-based hydrophilic interaction liquid chromatography (HILIC) particles were purchased from Sigma Aldrich (USA), whereas the protein standard marker and analytical-grade trypsin were obtained from Agilent Technologies (Santa Clara, CA, USA).

2.3. Venom protein extraction and pellet solubilization

The venom proteins were extracted following a method described previously [25]. Two milligrams (dry-weight) of the lyophilized crude venom (from each species) were solubilized in $50\ \mu\text{L}\ 1\times\ \text{PBS}$ (pH 7.4), vortexed for 10 min, and centrifuged at $15,700\times g$ for 5 min at 4°C . Acetone precipitation was performed on the supernatant by the addition of $600\ \mu\text{L}$ cold acetone, and the samples were incubated at -20°C for 15 min. Precipitated samples were centrifuged at $15,700\times g$ for 15 min at 4°C . Protein pellets were air-dried and resuspended in $100\ \mu\text{L}\ 1\times\ \text{PBS}$ buffer (pH 7.4). Protein concentration of each sample was quantified using the RC DC Protein Assay Kit 11 (Bio-Rad Laboratories) and confirmed with the microvolume protein concentration determination method. Data was acquired on a Nanodrop Spectrophotometer 2000c (Thermo Fisher Scientific, USA).

2.4. One-dimensional SDS-PAGE

Aliquots of each sample ($15\ \mu\text{g}$) were subjected to 12% SDS-PAGE under non-reducing conditions and ran for 90 min at 100 V. The gels were stained with Coomassie Brilliant Blue (G-250), and de-stained in 10% glacial acetic acid containing 1% glycerol. The resolved protein bands were visualized using a Molecular Imager PhorosFX Plus System (Bio-Rad, California, USA).

2.5. Hydrophilic interaction liquid chromatography (HILIC) and trypsin digestion

HILIC was carried out as described previously [25]. Fifty micrograms ($50\ \mu\text{g}$) of protein from each venom sample was suspended in a final concentration of 50 mM triethylammonium bicarbonate (TEAB; Sigma). The proteins were reduced with a final concentration of 10 mM Dithiothreitol (DTT; Sigma) in 50 mM TEAB for 40 min at 56°C . Samples were cooled to room temperature and alkylated with 30 mM iodoacetamide (Sigma) in 50 mM TEAB at 25°C and the mixture was kept in the dark for 30 min. After alkylation, two-fold dilutions of the samples were made with a binding buffer (100 mM ammonium acetate, 30% acetonitrile, pH 4.5). The protein solution was added to pre-equilibrated MagResyn® HILIC magnetic particles (Resyn Biosciences) prepared according to the manufacturer’s instructions and incubated overnight at 4°C . After binding, the supernatant was removed and the magnetic particles were washed twice with 95% acetonitrile for 1 min. Thereafter, the magnetic particles were suspended in 50 mM ammonium formate (pH 8.0) containing trypsin (Promega) to a final enzyme: protein ratio of 1:10 and the content was incubated overnight at 27°C with constant shaking. Following the digestion, peptides were recovered with 1% trifluoroacetic acid (TFA); which were then incubated at room temperature for 3 min and analyzed using liquid chromatography with tandem mass spectrometry (LC-MS/MS).




Fig. 2. Relative distribution of protein families in the venom proteome of *B. arietans*.

2.6. Characterization of venom proteins

Liquid chromatography was performed as described by Lomonte et al. [26] with slight modifications using an UltiMate™ 3000 RSLCnano System (Thermo Fisher Scientific, USA) equipped with a C₁₈ trap column (5 mm × 300 μm; Thermo Fisher Scientific) and a CSH C₁₈ analytical column (1.7 μm, 25 cm × 75 μm), with a linear gradient of 0.1% trifluoroacetic acid in water (solution A) and acetonitrile (solution B). The samples were loaded on the trap column and the flow rate was set at 250 nL/min and the gradient was generated as follows: 5–35% solution B for 60 min and 35–50% solution B for 60–75 min. Chromatography was performed at 40 °C, and the outflow was delivered to the mass spectrometer through a stainless-steel nano-bore emitter. LC-MS analysis was performed on a Fusion mass spectrometer (Thermo Scientific, San Jose, CA) equipped with a nanospray ion source (Nanospray Flex™ Ion Sources, Thermo Fisher Scientific) coupled to a Dionex Ultimate 3000 RSLC nano-HPLC system. Peptides recovered from the on-bead HILIC digestion for each venom sample was introduced through a stainless-steel emitter. Data were collected in positive mode with a spray voltage and ion transfer capillary set at 1.8 kV and 280 °C respectively. Spectra were internally calibrated using polysiloxane ions at *m/z* 445.12003 and 371.10024. The first MS scan was performed using the orbitrap detector at 120,000 resolutions over an *m/z* range of 350–1650 with an AGC target at 3E5 and maximum injection time of 40 ms. Data were acquired in profile mode. The second MS scan was performed using monoisotopic precursor selection for the ion with charges of +2 to +7 with error tolerance set at ±10 ppm. Precursor ions were excluded for 60 s after fragmentation. Precursor ions were selected for fragmentation

in the HCD mode using the quadrupole mass analyzer with HCD energy set to 30%. Fragment ions were detected in the orbitrap mass analyzer at a resolution of 30000. The AGC target was set to 5E4 and the maximum injection time was set to 80 ms. Data were acquired in centroid mode.

2.7. Data analysis

The mass spectra obtained were subjected to analysis and processing using Proteome Discoverer v1.4 software (Thermo Fisher Scientific, USA) and the Sequest and Amanda algorithms, respectively. Database interrogation was performed against a concatenated database created by concatenating all 'snake protein' entries, including the previously analyzed transcriptome data of the two snakes in the Uniprot-Serpentes database with the cRAP contaminant database (<https://www.thegpm.org/crap/>). The precursor mass tolerance was set to 10 ppm and fragment mass tolerance was set to 0.02 Da. Peptide validation was performed using the Target-Decoy PSM validator node. The search results were subjected to Scaffold Q+ version 4.10.0 for further validation (www.proteomesoftware.com). Peptide identification, by peptide-spectrum match approach against UniProt-Serpentes database, and subsequent assembly, matching, and identification of protein sequences were performed using X! Tandem and Sequest search engines. The search engines were incorporated into Scaffold Q+, version 4.10.0 at 99% protein threshold, 95% peptide threshold, and two-peptide minimum criterion. The relative abundances (expressed as percentage of the total venom proteins) of the protein families were obtained by the unique spectral count strategy for label-free quantitative proteomics [27]. For accurate quantitation, Abacus computational tool for spectral




Fig. 3. Relative distribution of protein families in the venom proteome of *E. ocellatus*.

counts was used to implement a distributed Normalized Spectral Abundance Factor (dNSAF), to accurately account for peptides shared across multiple proteins [28,29]. The mass spectrometric data has been deposited into the ProteomeXchange Consortium via the PRIDE [30] with an identifier PXD024638.

3. Results and discussion

3.1. One-dimensional SDS-PAGE of the crude snake venoms

A fraction (15 µg) of each crude venom was size-fractionated on a one-dimensional SDS polyacrylamide gel and visualized using Coomassie Brilliant Blue R-250 dye (Fig. 1). The protein bands observed for the two venoms were distinctly different and clearly separated within the molecular weight ranges of 7 kDa, 22 kDa and 100 kDa (Fig. 1). The distinct pattern on the 1D gel correlated with the LC-MS/MS analysis which showed that *B. arietans* had serine proteases and metalloproteinases as the most abundant venom proteins (Fig. 2), while the venom of *E. ocellatus* was composed of three major protein families; i.e. metalloproteinases, phospholipase A₂s, and serine proteases (Fig. 3).

3.2. Snake venom proteomic characterization using LC-MS/MS

Annotation was against the comprehensive UniProt Serpentes database where all curated data are available, including the venom gland transcriptome dataset of Nigerian *E. ocellatus* and *B. arietans*. A total of

79 proteins were detected in the venom of *B. arietans* whereas 85 proteins were detected in the venom of *E. ocellatus* (Table 1) when the digested crude venom was analyzed using LC-MS/MS. Table 2 summarizes the difference and similarities of the protein and protein families. These vipers shared a total of 78 proteins and 14 protein families in common. In addition, *E. ocellatus* had 4 protein families namely; phospholipase A₂ inhibitors, snake venom glutaminyl cyclases, secretory phospholipase A₂ receptors and globin exclusive to its venom (Table 2). Overall, the protein similarity index was found to be 91.8%. Thus, all the proteins detected (in the molecular mass range of 4–93 kDa) (Table 3) belonged to 14 protein families in *B. arietans* and 18 protein families in *E. ocellatus* (Table 2). The most abundant proteins in the venom of *B. arietans* were serine proteases (22%) and metalloproteases (21%) (Fig. 2), followed by C-type lectins/snaclecs (10.65%), phospholipase A₂ (10.6%), L-amino acid oxidases (8.72%) and vascular endothelial growth factors (8.14%). Three-finger toxins (3.35%), disintegrins (3.4%), 5'-nucleotidases (2.5%), venom nerve growth factor (2.16%) and cysteine-rich venom proteins (2.05%) were found in relatively low abundance. Kunitz-type peptides, peroxiredoxin and phospholipase B were the least abundant protein families (<1%) identified in *B. arietans* (Fig. 2). On the other hand, the most abundant proteins identified in the venom of *E. ocellatus* include metalloproteinases (34.84%), phospholipase A₂ (21.19%), and serine proteases (15.05%) (Fig. 3). C-type lectins/snaclecs (3.95%), three-finger toxins (2.16%), vascular endothelial growth factors (2.67%), cysteine-rich venom proteins (2.9%) and phospholipase A₂ inhibitors (2.64%) were found in much lower

Table 1
Protein identification in the venoms of *B. arietans* and *E. ocellatus*.

Protein/peptide detected	Originating species	Originating species	Originating species
S/ no.	<i>Echis ocellatus</i>	<i>Bitis arietans</i>	
1	Acidic phospholipase A ₂	<i>Echis ocellatus</i>	Acidic phospholipase A ₂
2	Acidic phospholipase A ₂	<i>Naja mossambica</i>	Acidic phospholipase A ₂
3	Alpha-fibrinogenase	<i>Macrovipera lebetina</i>	Alpha-fibrinogenase
4	Alpha-fibrinogenase-like	<i>Daboia siamensis</i>	Alpha-fibrinogenase-like
5	Beta-fibrinogenase brevinase	<i>Gloydius blomhoffii</i>	Beta-fibrinogenase brevinase
6	C-type lectin 1	<i>Bitis gabonica</i>	C-type lectin 1
7	Cationic trypsin	<i>Bos taurus</i>	Cationic trypsin
8	Coagulation factor X-activating enzyme heavy chain	<i>Daboia siamensis</i>	Coagulation factor X-activating enzyme heavy chain
9	Coagulation factor X-activating enzyme heavy chain	<i>Macrovipera lebetina</i>	Coagulation factor X-activating enzyme heavy chain
10	Cysteine-rich venom protein	<i>Trimeresurus stejnegeri</i>	Cysteine-rich venom protein
11	Cytotoxin 1	<i>Naja mossambica</i>	Cytotoxin 1
12	Disintegrin EO5A	<i>Echis ocellatus</i>	Disintegrin EO5A
13	Disintegrin bitistatin	<i>Bitis arietans</i>	Disintegrin bitistatin
14	Glutaminy-peptide cyclotransferase	<i>Boiga dendrophila</i>	Kunitz-type serine protease inhibitor bitisilin-1
15	Hemoglobin subunit beta	<i>Erythrolamprus mliaris</i>	Kunitz-type serine protease inhibitor bitisilin-3
16	Kunitz-type serine protease inhibitor bitisilin-1	<i>Bitis gabonica</i>	L-amino-acid oxidase
17	Kunitz-type serine protease inhibitor bitisilin-3	<i>Bitis gabonica</i>	L-amino-acid oxidase
18	L-amino-acid oxidase	<i>Bitis gabonica</i>	L-amino-acid oxidase
19	L-amino-acid oxidase	<i>Echis ocellatus</i>	Long neurotoxin 1
20	L-amino-acid oxidase	<i>Pseudechis australis</i>	Peroxiredoxin-4
21	Long neurotoxin 1	<i>Naja anchietae</i>	Phospholipase A ₂ homolog
22	Peroxiredoxin-4	<i>Crotalus atrox</i>	Phospholipase-B 81
23	Phospholipase A ₂ homolog	<i>Echis ocellatus</i>	Serine protease harobin
24	Phospholipase A ₂ inhibitor 1	<i>Protobothrops flavoviridis</i>	Serine protease sp-Eoc49
25	Phospholipase-B 81	<i>Drysdalia coronoides</i>	Short neurotoxin 1
26	Secretory phospholipase A ₂ receptor	<i>Pongo abelii</i>	Snaclec 2
27	Serine protease harobin	<i>Hydrophis hardwickii</i>	Snaclec 3
28	Serine protease sp-Eoc49	<i>Echis ocellatus</i>	Snaclec agglucetin subunit beta-1
29	Short neurotoxin 1	<i>Naja nivea</i>	Snaclec bitiscetin subunit alpha
30	Snaclec 2	<i>Bitis gabonica</i>	Snaclec bitiscetin subunit beta
31	Snaclec 3	<i>Bitis gabonica</i>	Snaclec clone 2100755
32	Snaclec CTL-Eoc124	<i>Echis ocellatus</i>	Snake venom 5'-nucleotidase
33	Snaclec CTL-Eoc125	<i>Echis ocellatus</i>	Snake venom metalloproteinase ACLH
34	Snaclec agglucetin subunit beta-1	<i>Deinagkistrodon acutus</i>	Snake venom metalloproteinase kistomin
35	Snaclec bitiscetin subunit alpha	<i>Bitis arietans</i>	Snake venom metalloproteinase lebetase-4
36	Snaclec bitiscetin subunit beta	<i>Bitis arietans</i>	Snake venom metalloproteinase-disintegrin-like mocarhagin
37	Snaclec clone 2100755	<i>Deinagkistrodon acutus</i>	Snake venom serine protease BthaTL
38	Snaclec convulxin subunit beta	<i>Crotalus durissus terrificus</i>	Snake venom serine protease HS114
39	Snake venom 5'-nucleotidase	<i>Crotalus adamanteus</i>	Snake venom serine protease NaSP
40	Snake venom metalloproteinase ACLH	<i>Agkistrodon contortrix laticinctus</i>	Snake venom serine protease
41	Snake venom metalloproteinase kistomin	<i>Calloselasma rhodostoma</i>	Snake venom vascular endothelial growth factor toxin barietin
42	Snake venom metalloproteinase lebetase-4	<i>Macrovipera lebetina</i>	Thrombin-like enzyme cerastocytin
43	Snake venom metalloproteinase-disintegrin-like mocarhagin	<i>Naja mossambica</i>	Vascular endothelial growth factor A
44	Snake venom serine protease BthaTL	<i>Bothrops alternatus</i>	Venom nerve growth factor 2
45	Snake venom serine protease HS114	<i>Bothrops jararaca</i>	Weak neurotoxin 7
46	Snake venom serine protease NaSP	<i>Naja atra</i>	Weak toxin CM-2a
47	Snake venom serine protease	<i>Philodryas olfersii</i>	Zinc metalloproteinase homolog-disintegrin albolatin
48	Snake venom vascular endothelial growth factor toxin barietin	<i>Bitis arietans</i>	Zinc metalloproteinase leucurolysin-B
49	Thrombin-like enzyme cerastocytin	<i>Cerastes cerastes</i>	Zinc metalloproteinase-disintegrin BA-5A
50	Venom nerve growth factor 2	<i>Naja sputatrix</i>	Zinc metalloproteinase-disintegrin B1atH1
51	Weak neurotoxin 7	<i>Naja naja</i>	Zinc metalloproteinase-disintegrin bilitoxin-1
52	Weak toxin CM-2a	<i>Naja annulifera</i>	Zinc metalloproteinase-disintegrin-like BfMP
53	Zinc metalloproteinase homolog-disintegrin albolatin	<i>Trimeresurus albolabris</i>	Zinc metalloproteinase-disintegrin-like EoMP06
54	Zinc metalloproteinase leucurolysin-B	<i>Bothrops leucurus</i>	Zinc metalloproteinase-disintegrin-like EoVMP2
55	Zinc metalloproteinase-disintegrin BA-5A	<i>Bitis arietans</i>	Zinc metalloproteinase-disintegrin-like Eoc1
56	Zinc metalloproteinase-disintegrin B1atH1	<i>Bothriechis lateralis</i>	Zinc metalloproteinase-disintegrin-like HF3
57	Zinc metalloproteinase-disintegrin bilitoxin-1	<i>Agkistrodon bilineatus</i>	Zinc metalloproteinase-disintegrin-like MTP9
58	Zinc metalloproteinase-disintegrin-like BfMP	<i>Bungarus fasciatus</i>	Zinc metalloproteinase-disintegrin-like NaMP
59	Zinc metalloproteinase-disintegrin-like EoMP06	<i>Echis ocellatus</i>	Zinc metalloproteinase-disintegrin-like
60	Zinc metalloproteinase-disintegrin-like EoVMP2	<i>Echis ocellatus</i>	Zinc metalloproteinase-disintegrin-like TSV-DM
61	Zinc metalloproteinase-disintegrin-like Eoc1	<i>Echis ocellatus</i>	Zinc metalloproteinase-disintegrin-like VLAIP-A
62	Zinc metalloproteinase-disintegrin-like HF3	<i>Bothrops jararaca</i>	Zinc metalloproteinase-disintegrin-like VLAIP-B
63	Zinc metalloproteinase-disintegrin-like MTP9	<i>Drysdalia coronoides</i>	Zinc metalloproteinase-disintegrin-like acurhagin
64	Zinc metalloproteinase-disintegrin-like NaMP	<i>Naja atra</i>	Zinc metalloproteinase-disintegrin-like atrolysin-A

(continued on next page)

Table 1 (continued)

Protein/peptide detected	Originating species	Originating species	Originating species
S/ no.	<i>Echis ocellatus</i>	<i>Bitis arietans</i>	
65	Zinc metalloproteinase-disintegrin-like	<i>Cerberus rynchops</i>	Zinc metalloproteinase-disintegrin-like batroxstatin-1
66	Zinc metalloproteinase-disintegrin-like TSV-DM	<i>Trimeresurus stejnegeri</i>	Zinc metalloproteinase-disintegrin-like batroxstatin-3
67	Zinc metalloproteinase-disintegrin-like VLAIP-A	<i>Macrovipera lebetina</i>	Zinc metalloproteinase-disintegrin-like berythracivase
68	Zinc metalloproteinase-disintegrin-like VLAIP-B	<i>Macrovipera lebetina</i>	Zinc metalloproteinase-disintegrin-like bothrojarin-2
69	Zinc metalloproteinase-disintegrin-like acurhagin	<i>Deinagkistrodon acutus</i>	Zinc metalloproteinase-disintegrin-like brevilysin H2a
70	Zinc metalloproteinase-disintegrin-like atrolysin-A	<i>Crotalus atrox</i>	Zinc metalloproteinase-disintegrin-like cobrin
71	Zinc metalloproteinase-disintegrin-like batroxstatin-1	<i>Bothrops atrox</i>	Zinc metalloproteinase-disintegrin-like daborhagin-K
72	Zinc metalloproteinase-disintegrin-like batroxstatin-3	<i>Bothrops atrox</i>	Zinc metalloproteinase-disintegrin-like stejnihagin-A
73	Zinc metalloproteinase-disintegrin-like berythracivase	<i>Bothrops erythromelas</i>	Zinc metalloproteinase-disintegrin-like stejnihagin-B
74	Zinc metalloproteinase-disintegrin-like bothrojarin-2	<i>Bothrops jararaca</i>	Zinc metalloproteinase/disintegrin
75	Zinc metalloproteinase-disintegrin-like brevilysin H2a	<i>Gloydius brevicaudus</i>	Zinc metalloproteinase/disintegrin
76	Zinc metalloproteinase-disintegrin-like cobrin	<i>Naja kaouthia</i>	Zinc metalloproteinase/disintegrin
77	Zinc metalloproteinase-disintegrin-like daborhagin-K	<i>Daboia russelii</i>	Metalloproteinase/disintegrin
78	Zinc metalloproteinase-disintegrin-like stejnihagin-A	<i>Trimeresurus stejnegeri</i>	Zinc metalloproteinase/disintegrin
79	Zinc metalloproteinase-disintegrin-like stejnihagin-B	<i>Trimeresurus stejnegeri</i>	Zinc metalloproteinase/disintegrin PMMP-1
80	Zinc metalloproteinase/disintegrin	<i>Protobothrops mucrosquamatus</i>	
81	Zinc metalloproteinase/disintegrin	<i>Trimeresurus gramineus</i>	
82	Zinc metalloproteinase/disintegrin	<i>Calloselasma rhodostoma</i>	
83	Metalloproteinase/disintegrin	<i>Echis ocellatus</i>	
84	Zinc metalloproteinase/disintegrin	<i>Macrovipera lebetina</i>	
85	Zinc metalloproteinase/disintegrin PMMP-1	<i>Protobothrops mucrosquamatus</i>	

Proteins were identified using Multi-Dimensional Protein Identification Technology (MuDPIT) incorporated on Scaffold Proteome software version 4.10.0 at 99% protein threshold, 95% peptide threshold, 0.5% false discovery rate (FDR), and two-peptide minimum criterion. Peptides and proteins were searched against the UniprotKB- Serpentes-database.

abundance. Disintegrins, 5'-nucleotidases, L-amino acid oxidases, snake venom glutaminyl cyclases, secretory phospholipase A₂ receptors and Kunitz-type peptides families were very low in abundance (<3%), while phospholipase B, globin and venom nerve growth factor were virtually absent with percentage abundance of <1%, as shown in Fig. 3.

3.3. Relative distribution of the protein families

Increase in the accuracy of spectral count quantitation emerged following the advancements in bioinformatics tools [31]. As a general rule in spectral count, the number of spectra associated with a peptide shared by different proteins is used for computing spectral count of each protein with similar peptide. This has a limitation of introducing inaccuracy in the estimation of protein abundances, in that if one or more peptides are shared between high and low-abundant proteins, the proportion of the low-abundant protein tends to be overestimated [32]. This limitation was circumvented by implementing the distributed Normalized Spectral Abundance Factor (dNSAF), where shared spectrum count is distributed based on the relative proportions of unique spectra assigned to the proteins containing the shared peptides [28]. Thus, Abacus software was used so as to extract and adjust the spectral counts from MS/MS data sets to accurately account for peptides shared across multiple proteins, through normalization steps [29]. In principle, the composition of the analyzed venoms indicated that metalloproteinases (SVMPs), serine proteinases (SVSPs), and phospholipases A₂ (PLA₂S) constituted the major proportions of the venom proteomes, which is consistent with the report of previous studies on viper venomics [16,33,

34]. However, a deep insight into the proportion of these major protein families revealed a significant variation among species from different regions of the world; the protein composition of the venoms from vipers indigenous to Gaboon in West Africa showed that the percentages of PLA₂S, SVSPs and SVMPs in *B. arietans* were 4.3%, 19.5% and 38.5% respectively [34]. More so, PLA₂S, and SVMPs were the major toxin families, with varied proportions in the venoms of *Echis* species [35]. In the venoms of these vipers, indigenous to Nigeria, the percentages were higher; with PLA₂S, SVSPs and SVMPs having 10.6%, 22.31%, and 21.06% respectively in *B. arietans* while in *E. ocellatus*, 21.19%, 15.50% and 34.84% were recorded in PLA₂S, SVSPs and SVMPs respectively (Figs. 2 and 3). These variations are attributed to factors such as snake's gender, age [20], geographical origin [15–18] and diet [19]. It is believed that mutations in the venom-related genes may affect the venom composition [25]. Fifteen secondary protein families (Table 4) were detected, namely, kunitz-type peptides, cysteine-rich venom proteins, three-finger toxins, disintegrins, L-amino acid oxidases, C-type lectins/snaclecs, 5'-nucleotidases, vascular endothelial growth factors, phospholipase B, peroxiredoxin, phospholipase A₂ inhibitors, snake venom glutaminyl cyclases, globin, secretory phospholipase A₂ receptors, and venom nerve growth factors, with a distribution range between ~1% and 4% (Figs. 2 and 3).

Casewell et al. [22] showed a transcriptomic analysis of venom gland where snake venom metalloproteinases and serine proteases were the major toxin genes transcribed in Nigerian *B. arietans*, and were also found to be the most abundant toxin family secreted in the venom [22]. This is in tandem with our findings that SVSPs and SVMPs were the

Table 2

Summary of the proteins identified, presenting the similarities and differences of proteins and protein families.

Protein family	Number of proteins		
	<i>B. arietans</i>	<i>E. ocellatus</i>	Common to both venoms
Snake venom metalloproteinases	0	0	37
Snake venom Serine proteases	0	0	13
Phospholipase A _{2s}	0	0	3
Kunitz peptides	0	0	2
Cysteine-rich venom protein	0	0	1
Disintegrins	0	0	2
L-amino acid oxidases	0	0	3
Three-finger toxins	0	0	5
C-type lectins/Snaclects	0	3	7
5' Nucleotidases	0	0	1
Vascular endothelial growth factors	1	0	1
Phospholipase B	0	0	1
Peroxiredoxin	0	0	1
Phospholipase A ₂ inhibitor	0	1	0
Snake venom glutaminyl cyclases	0	1	0
Globin	0	1	0
Secretory phospholipase A ₂ receptor	0	1	0
Venom nerve growth factor	0	0	1
Total	1	7	78

The proteins identified were classified into 14 and 18 protein families in *B. arietans* and *E. ocellatus* respectively, with 78 proteins common to both venoms. There were 1 and 7 protein(s) exclusive to the venom of *B. arietans* and *E. ocellatus* respectively. *E. ocellatus* had additional 4 protein families (Phospholipase A₂ inhibitor, Snake venom glutaminyl cyclases, Globin and Secretory phospholipase A₂ receptor) exclusive to its venom.

dominant proteins in the venom of *B. arietans*. Similarly, snake venom metalloproteinases, phospholipase A₂ and serine proteases constituted the most abundant toxin genes transcribed in the venom glands of *E. ocellatus* from Nigeria [23]. The abundance of these 3 toxin families (PLA_{2s}, SVSPs and SVMPs) in the venom gland transcriptomic analyses correlates with the findings of this study. In a parallel comparison, the transcriptomic analyses showed that C-type lectins and L-amino oxidases were present in high abundances but in this study, they are present in low abundances 3.95% and 2.24% respectively in the venom proteome (Fig. 3). This disparity may be due to the fact that toxins are known to be transcribed at the highest level in the venom gland [22].

3.4. Pathological mechanisms of the toxin families

Snake venom toxins are thought to act synergistically in exerting a wide range of biochemical and toxicological effects [36,37]. Envenomation by the viper primarily gives rise to local effects and bleeding [38, 39]. Snake venom metalloproteinases (SVMPs) are zinc-dependent proteinase toxins that are responsible for many of the generally known pathological phenotypes in envenomation by the viper snake species [16,40–42] and play a key role in coagulopathies commonly associated with viper envenoming [40,41,43]. Similarly, SVMPs provoke a manifold of clinical manifestations, including hemorrhagic, pro-coagulant, anticoagulant, fibrinolytic, apoptotic, and antiplatelet activities [44, 45]. Thus, SVMPs are believed to evolve from ADAM (a Disintegrin and Metalloproteinase) proteins, precisely ADAM28, characterized by metalloproteinase, disintegrin-like and cysteine-rich domains [46], which induce hemorrhagic activity by rupturing the capillary vessels, resulting to extravasation in envenomed victims [47]. This occurs through cleavage of the basement membrane and adhesion proteins of the endothelial cells-matrix, causing the endothelial cells to detach and become thin, leading to the obstruction of the capillary walls and blood effusion [40,48]. More so, SVMPs alters homeostasis by disrupting

coagulation, through modulation of fibrinogenase and fibrolase that mediate the coagulation cascade which aids in eliciting their hemorrhagic role [49,50]. The venom of the viper snakes contained the three related structural classes of SVMPs, protein-type I, II and III. The P-I class has only a metalloproteinase domain, P-II class consists of a metalloproteinase and disintegrin domain, whereas the P-III class has a metalloproteinase domain, a disintegrin-like domain and a cysteine-rich domain. Among the protein-type I snake venom metalloproteinases, kistomin impairs platelet function by binding and cleaving of both platelet membrane glycoprotein Ib and glycoprotein VI [51]. ACLH exerts fibrinogenolytic and fibrinolytic effects [52], acurhagin preferentially cleaves Alpha chain of fibrinogen, followed by Beta chain, with minimal effects on the gamma chains [53]. Bilitoxin-1 represents protein-type II hemorrhagic toxin isolated from the venom of these snakes [54]. The catalytic activity of protein-type III hemorrhagic leucurolysin-B on extracellular matrix proteins leads to loss of capillary integrity, resulting in hemorrhage with alterations in platelet function [55]. Atrolysin A is a potent inhibitor of platelet aggregation, the region of the disintegrin-like domain is primarily responsible for blocking platelet aggregation [56]. Berythactivase contains metalloproteinase, disintegrin-like and cysteine-rich domains. Structurally, berythactivase is similar to snake-venom haemorrhagic metalloproteinases and functionally similar to group A prothrombin activators [57].

Snake venom serine proteinases (SVSPs) are multifunctional enzymes that induce toxicity in envenomed victims by affecting the hemostatic system. They alter blood coagulation processes, fibrinolysis and platelet aggregation [35,36,58].

In this study, the venom of *E. ocellatus* was found to be dominated by Phospholipases A₂ (PLA_{2s}) which constituted 21% of the venom proteome. Our data correlates with the previous studies in which PLA_{2s} were described as one of the commonest toxins in the venom of the front-fanged snakes [59,60]. Although PLA_{2s} were present in both venoms, it was found to be lower (10.6%) in the venom of *B. arietans* (Fig. 2). These toxins had molecular masses of 13 and 16 kDa (Table 3) and are important cause of myotoxic effect of viper envenomation [61,62], which often leads to severe necrosis [63]. In addition, PLA_{2s} from venoms of *Viperidae* snake family induce membranotropic effects [64, 65].

Other toxin families identified in the venom of these vipers were three-finger toxins, kunitz-type peptides, cysteine-rich secretory proteins, disintegrins, L-amino acid oxidases, C-type lectins, 5-nucleotidase, vascular endothelial growth factors and phospholipase B. These toxins were detected, albeit, at lower abundances in the venom of *E. ocellatus* (Fig. 3) compared to that of *B. arietans* (Fig. 2). The toxin 5'-nucleotidases inhibit the aggregation of platelet through the release of adenosine from adenosine monophosphate which binds to receptors on the platelets [66], this shows that they act in synergy with hemorrhagic toxins to elicit anticoagulant effect observed in snake bite envenoming [67,68].

L-amino acid oxidases (LAAOs) in snake venoms catalyze oxidative deamination of amino acids, leading to the release of hydrogen peroxide [69], which provokes toxicity by impacting on platelet aggregation and inducing hemorrhage that ultimately leads to apoptosis of vascular endothelial cells [70,71].

Additionally, the cysteine-rich venom proteins (CRISPs) are toxin families whose exact function in snake venoms is not known, they are however, potential blockers of Ca²⁺ and cyclic nucleotide-gated channels [72]. The C-type lectins/snaclecs (CTLs) are calcium-dependent non-enzymatic proteins that bind carbohydrates, mainly galactose [37] and play important roles in the agglutination of erythrocytes and aggregation of platelet [73].

Also, three-finger toxins (3FTXs) are non-enzymatic three-finger fold structure that is both neurotoxic and cytotoxic in nature [74]. 3FTXs are commonly found in the venoms of snakes from the Elapids family [16], they bind at the postsynaptic neuromuscular junctions, thereby inducing flaccid paralysis in victims [75]. Their biological activity in the venoms of *Viperidae* snake family needs to be explored [64]. However, their

Table 3
 Characteristics of the proteins identified in the venom extracts of *B. arietans* and *E. ocellatus*.

S/ No.	Protein/Toxin detected	Protein accession number	Mol. mass (kDa)	B. arietans				E. ocellatus			
				AA (%)	NET	TUSC	EUP	AA (%)	NET	TUSC	EUP
1	Acidic phospholipase A2	PA2A5_ECHOC	16	83	32	12	8	61	439	79	38
2	Acidic phospholipase A2	PA2A1_NAJMO	13	39	19	1	1	39	9	1	1
3	Alpha-fibrinogenase	VSPA_MACLB	29	8.5	29	0	0	8.1	20	0	0
4	Alpha-fibrinogenase-like	VSPAF_DABSI	28	3.5	28	0	0	0.8	20	1	1
5	Beta-fibrinogenase brevinase	VSPB_GLOBL	26	8.6	54	0	0	16	76	2	1
6	C-type lectin 1	LEC1_BITGA	19	16	4	2	2	8.2	2	1	1
7	Cationic trypsin	TRY1_BOVIN	26	16	5	3	3	8.5	2	1	1
8	Coagulation factor X-activating enzyme heavy chain	VM3CX_DABSI	70	8.1	33	0	0	21	43	9	4
9	Coagulation factor X-activating enzyme heavy chain	VM3CX_MACLB	69	8.5	14	0	0	18	54	4	3
10	Cysteine-rich venom protein	CRVP_TRIST	26	4.3	1	0	0	13	14	0	0
11	Cytotoxin 1	3SA1_NAJMO	7	47	9	0	0	33	30	0	0
12	Disintegrin EO5A	DID5A_ECHOC	12	9.6	7	0	0	36	31	0	0
13	Disintegrin bitistatin	VM2_BITAR	9	88	80	11	6	27	6	12	6
14	Glutaminyl-peptide cyclotransferase	QPCT_BOIDE	42	0.0	0	0	0	17	12	1	1
15	Hemoglobin subunit beta	HBB_ERYML	16	0.0	0	0	0	6.2	3	1	1
16	Kunitz-type serine protease inhibitor bitsilin-1	VKT1_BITGA	10	18	5	2	2	10	2	2	2
17	Kunitz-type serine protease inhibitor bitsilin-3	VKT3_BITGA	17	42	12	5	4	17	7	2	2
18	L-amino-acid oxidase	OXLA_BITGA	7	12	4	0	0	20	7	1	1
19	L-amino-acid oxidase	OXLA_ECHOC	57	34	58	1	1	63	109	1	1
20	L-amino-acid oxidase	OXLA_PSEAU	59	1.4	2	0	0	6.2	8	0	0
21	Long neurotoxin 1	3L21_NAJAC	8	90	60	6	2	93	64	11	5
22	Peroxiredoxin-4	PRDX4_CROAT	4	58	4	2	2	31	2	1	1
23	Phospholipase A2 homolog	PA2HS_ECHOC	16	60	42	19	13	78	312	75	46
24	Phospholipase A2 inhibitor 1	PLI1_PROFL	22	0.0	0	0	0	8.5	20	1	1
25	Phospholipase-B 81	PLB_DRYCN	64	19	121	14	11	20	73	14	11
26	Secretory phospholipase A2 receptor	PLA2R_PONAB	18	0.0	0	0	0	1.2	3	1	1
27	Serine protease harobin	VSPHA_HYDHA	29	7.9	21	3	1	7.9	21	4	3
28	Serine protease sp-Eoc49	VSP_ECHOC	28	24	27	4	4	49	51	18	13
29	Short neurotoxin 1	3S11_NAJNI	7	61	8	0	0	66	17	2	2
30	Snaclec 2	SL2_BITGA	18	35	26	6	5	31	18	7	6
31	Snaclec 3	SL3_BITGA	18	18	7	3	2	24	9	4	2
32	Snaclec CTL-Eoc124	SL124_ECHOC	17	0.0	0	0	0	65	33	14	9
33	Snaclec CTL-Eoc125	SL125_ECHOC	18	0.0	0	0	0	47	11	6	6
34	Snaclec agglucetin subunit beta-1	SLB1_DEIAC	17	9.6	2	0	0	9.6	4	0	0
35	Snaclec bitiscetin subunit alpha	SLA_BITAR	15	96	124	0	0	91	82	0	0
36	Snaclec bitiscetin subunit beta	SLB_BITAR	15	60	71	27	13	50	60	18	7
37	Snaclec clone 2100755	SL_DEIAC	18	4.5	4	27	13	14	5	18	7
38	Snaclec convulxin subunit beta	SLB_CRODU	17	0.0	0	0	0	11	4	0	0
39	Snake venom 5'-nucleotidase	V5NTD_CROAD	65	39	43	11	5	30	52	5	3
40	Snake venom metalloproteinase ACLH	VM1AH_AGKCL	46	5.2	3	0	0	7.1	7	0	0
41	Snake venom metalloproteinase kistomin	VM1K_CALRH	47	5.3	10	0	0	2.9	4	0	0
42	Snake venom metalloproteinase lebetase-4	VM1L4_MACLB	24	12	4	0	0	25	17	1	1
43	Snake venom metalloproteinase-disintegrin-like mocarhagin	VM3M1_NAJMO	68	2.0	30	4	2	4.4	18	5	2
44	Snake venom serine protease BthaTL	VSPTL_BOTAL	26	5.6	7	0	0	5.6	6	0	0
45	Snake venom serine protease HS114	VSP14_BOTJA	28	5.4	1	0	0	5.4	3	0	0
46	Snake venom serine protease NaSP	VSP1_NAJAT	31	7.4	3	0	0	7.4	2	0	0
47	Snake venom serine protease	VSP_PHIOL	28	15	6	1	1	14	3	1	1
48	Snake venom vascular endothelial growth factor toxin barietin	TXVE_BITAR	17	55	302	30	15	59	107	24	13
49	Thrombin-like enzyme cerastocytin	VSP_P_CERCE	28	13	67	4	3	13	48	3	3
50	Venom nerve growth factor 2	NGFV2_NAJSP	27	13	10	0	0	18	10	0	0
51	Weak neurotoxin 7	3NO27_NAJNA	8	5	2	1	1	12	9	0	0
52	Weak toxin CM-2a	3SOK2_NAJHA	7	46	1	2	1	46	3	3	2
53	Zinc metalloproteinase homolog-disintegrin albolatin	VM2AL_TRIAB	54	5.4	13	2	1	12	13	1	1
54	Zinc metalloproteinase leucurolysin-B	VM3LB_BOTLC	36	13	8	1	1	18	26	1	1
55	Zinc metalloproteinase-disintegrin BA-5A	VM25A_BITAR	59	13	10	6	5	31	96	28	21
56	Zinc metalloproteinase-disintegrin BlatH1	VM2H1_BOTLA	54	4.1	10	0	0	2.5	4	0	0
57	Zinc metalloproteinase-disintegrin bilitoxin-1	VM2B1_AGKBI	32	13	8	3	2	10	6	3	2
58	Zinc metalloproteinase-disintegrin-like BfMP	VM3_BUNFA	68	1.3	4	0	0	3.8	7	0	0
59	Zinc metalloproteinase-disintegrin-like EoMP06	VM3E6_ECHOC	58	17	21	6	4	92	165	28	17
60	Zinc metalloproteinase-disintegrin-like EoVMP2	VM3E2_ECHOC	69	19	20	6	5	55	241	117	66
61	Zinc metalloproteinase-disintegrin-like Eoc1	VM3E1_ECHOC	69	22	59	6	8	57	201	68	36
62	Zinc metalloproteinase-disintegrin-like HF3	VM3H3_BOTJA	68	9.6	14	0	0	13	30	0	0
63	Zinc metalloproteinase-disintegrin-like MTP9	VM39_DRYCN	68	1.3	4	0	0	5.4	8	0	0
64	Zinc metalloproteinase-disintegrin-like NaMP	VM3_NAJAT	69	3	5	1	1	1.6	2	1	1
65	Zinc metalloproteinase-disintegrin-like	VM3_CERRY	69	5.7	9	0	0	6.2	4	0	0
66	Zinc metalloproteinase-disintegrin-like TSV-DM	VM3TM_TRIST	69	8.2	12	0	0	6.8	7	0	0
67	Zinc metalloproteinase-disintegrin-like VLAIP-A	VM3VA_MACLB	69	19	58	7	6	22	92	10	7
68	Zinc metalloproteinase-disintegrin-like VLAIP-B	VM3VB_MACLB	69	14	37	2	2	11	38	2	1
69	Zinc metalloproteinase-disintegrin-like acurhagin	VM3AH_DEIAC	69	9.7	14	1	1	5.4	9	1	1

(continued on next page)

Table 3 (continued)

S/ No.	Protein/Toxin detected	Protein accession number	Mol. mass (kDa)	B. arietans				E. ocellatus			
				AA (%)	NET	TUSC	EUP	AA (%)	NET	TUSC	EUP
70	Zinc metalloproteinase-disintegrin-like atrolysin-A	VM3AA_CROAT	47	4.3	4	1	1	7.6	12	3	3
71	Zinc metalloproteinase-disintegrin-like batroxstatin-1	VM31_BOTAT	46	1.9	4	0	0	5.7	3	1	1
72	Zinc metalloproteinase-disintegrin-like batroxstatin-3	VM33_BOTAT	69	8	7	0	0	4.3	11	1	1
73	Zinc metalloproteinase-disintegrin-like berythracivase	VM3BE_BOTER	69	4.9	9	0	0	9	14	3	1
74	Zinc metalloproteinase-disintegrin-like bothrojarin-2	VM3B2_BOTJA	24	7.8	2	0	0	14	20	4	4
75	Zinc metalloproteinase-disintegrin-like brevilysin H2a	VM32A_GLOBR	47	6.4	6	0	0	11	29	0	0
76	Zinc metalloproteinase-disintegrin-like cobrin	VM3_NAJKA	68	4.7	2	1	1	8.2	5	2	2
77	Zinc metalloproteinase-disintegrin-like daborhagin-K	VM3DK_DABRR	70	7.3	5	2	2	15	34	5	5
78	Zinc metalloproteinase-disintegrin-like stejnihagin-A	VM3SA_TRIST	68	7.7	13	0	0	7.2	11	2	1
79	Zinc metalloproteinase-disintegrin-like stejnihagin-B	VM3SB_TRIST	68	7	11	0	0	8.3	13	0	0
80	Zinc metalloproteinase/disintegrin	VM2P3_PROMU	46	9.2	8	0	0	2.9	1	0	0
81	Zinc metalloproteinase/disintegrin	VM3G1_TRIGA	48	2.1	3	0	0	7.4	22	0	0
82	Zinc metalloproteinase/disintegrin	VM2RH_CALRH	54	12	16	0	0	6.3	13	2	1
83	Metalloproteinase/disintegrin	VM2OC_ECHOC	55	11	22	10	4	39	216	45	29
84	Zinc metalloproteinase/disintegrin	VM2L2_MACLB	53	13	17	0	0	17	2	4	2
85	Zinc metalloproteinase/disintegrin PMMP-1	VM2P1_PROMU	53	7.9	13	0	0	9.8	11	1	1

The characterization of proteins was achieved using Sequest and X!Tandem incorporated Scaffold Proteome Software 4.10.0: 2019/100.

Abbreviation: Mol. mass: molecular mass, %AA: percentage of amino acids, NET: Number of estimated enzymatic cleavage, EUP: Exclusive unique Peptide, TUPC: Total Unique Protein Count.

relative proportion was found to be 3.35% and 2.16% in the venoms of *B. arietans* (Fig. 2) and *E. ocellatus* (Fig. 3) respectively.

Disintegrins are venom toxins that were first detected in viper snakes [76], and are potent inhibitor of platelet aggregation, hence, antagonize the clotting of blood [77], thereby contributing to lethality in envenomed victims.

Moreover, snake venom vascular endothelial growth factors were distributed in minute quantities notably in the venom of *E. ocellatus*. They possess vasculotoxin-like activity which makes snakebite site liable to rapid infiltration of venom into victims [78]. In the case of kunitz-type peptides, a very low percentage (1%) in the venom of these vipers (Figs. 2 and 3) was obtained with molecular mass range of 10–17 kDa (Table 3). Phospholipase B (PLB) was among the least abundant proteins with 1.8% (Figs. 2) and 0.6% (Fig. 3) in the venoms of *B. arietans* and *E. ocellatus* respectively. In snake venomomics analysis, much interest has not been given to PLB probably because the enzyme is not commonly identified in snake venoms [79], therefore, there are limited data on this enzyme, and hence, the exact pathological function of PLB in snake venom has not been explained. Several studies reported the presence of PLB in some viper venoms [79,80], which is in accord with our findings.

4. Snake venom phospholipase A₂ inhibitor

It is known that phospholipase A₂ is one of the venom toxins that cause the disorder associated with snakebite envenoming. Hence, inhibiting the lethal actions of snake venom Phospholipase A₂ presents a strategic target for therapeutic interference [81]. Interestingly, Phospholipase A₂ inhibitor, a potential antitoxin with molecular mass of 22 kDa was detected in the venom of *E. ocellatus*. Available literature reports that the blood plasma of numerous snake species are naturally composed of endogenous phospholipase A₂ inhibitors, generally known as snake blood phospholipase A₂ inhibitors (sbPLIs). Primarily, they neutralize deleterious effects of phospholipase A₂ [82,83]. It implies that sbPLIs exert self-protection role against the enzymes of their own venom, which eventually reach the circulatory system [82,83]. What seems new is the fact that phospholipase A₂ inhibitory protein was detected in the venom, revealing that the protein is not only expressed in the snake blood serum. Exploring this promising protein may serve as

antidote to PLA₂-related disorders by giving rise to clinically relevant and specific antivenom that targets phospholipase A₂.

5. Conclusion

In addition to the detection of a venom-based antitoxin; venom phospholipase A₂ inhibitor, this study has provided a baseline information on the venom proteome of *E. ocellatus* and *B. arietans* indigenous to Nigeria, where *E. ocellatus* producing more of the toxins. In both snake venoms, their protein profile was found to be 91.8% similar and SVMPs, PLA₂s and SVSPs were the major toxin families. *E. ocellatus* had 34.84%, 21.19% and 15.50% while in *B. arietans*, 22.31%, 21.06% and 10.6% were recorded for SVSPs, SVMPs and PLA₂s respectively. Going by these findings, one can say that although the envenomation by these snakes may present some common pathogenic signs due to some similarities in the venom composition, their venoms are very much different when the toxins are taken into consideration and as such, this can serve as a guide in the use and design of a potential antivenom.

Authors' contributions

FAA, EJD, ABS: Conceptualization. ABS, AS, MSA: Supervision. EJD, FAA, GM, AK: Investigation. FAA, EJD: Data analysis. EJD: Writing original draft preparation. All authors critically reviewed and edited the final manuscript version.

Funding

This work did not receive any external funding from public or private sectors.

Declaration of competing interest

The authors declare no competing interests.

Data availability

The mass spectrometry/proteomics data have been deposited into

Table 4

Classification of the detected venom proteomes of *B. arietans* and *E. ocellatus* based on families.

Proteins	Protein families
Acidic phospholipase A ₂ Acidic phospholipase A ₂ Phospholipase A ₂ homolog	Phospholipase A ₂ s
Phospholipase-B 81	Phospholipase B
Alpha-fibrinogenase Alpha-fibrinogenase-like Beta-fibrinogenase brevinase Cationic trypsin Coagulation factor X-activating enzyme heavy chain Coagulation factor X-activating enzyme heavy chain Serine protease harobin Serine protease sp-Eoc49 Snake venom serine protease BthaTL Snake venom serine protease HS114 Snake venom serine protease NaSP Snake venom serine protease Thrombin-like enzyme cerastocytin	Snake venom serine proteases
Kunitz-type serine protease inhibitor bitisilin-1 Kunitz-type serine protease inhibitor bitisilin-3	Kunitz peptides
Snake venom metalloproteinase ACLH Snake venom metalloproteinase kistomin Snake venom metalloproteinase lebetase-4 Snake venom metalloproteinase-disintegrin-like mocarhagin Zinc metalloproteinase homolog-disintegrin albolatin Zinc metalloproteinase leucurolysin-B Zinc metalloproteinase-disintegrin BA-5A Zinc metalloproteinase-disintegrin Blath1 Zinc metalloproteinase-disintegrin bilitoxin-1 Zinc metalloproteinase-disintegrin-like BfMP Zinc metalloproteinase-disintegrin-like EoMPO6 Zinc metalloproteinase-disintegrin-like EoVMP2 Zinc metalloproteinase-disintegrin-like Eoc1 Zinc metalloproteinase-disintegrin-like HF3 Zinc metalloproteinase-disintegrin-like MTP9 Zinc metalloproteinase-disintegrin-like NaMP Zinc metalloproteinase-disintegrin-like Zinc metalloproteinase-disintegrin-like TSV-DM Zinc metalloproteinase-disintegrin-like VLAIP-A Zinc metalloproteinase-disintegrin-like VLAIP-B Zinc metalloproteinase-disintegrin-like acurhagin Zinc metalloproteinase-disintegrin-like atrolysin-A Zinc metalloproteinase-disintegrin-like batroxstatin-1 Zinc metalloproteinase-disintegrin-like batroxstatin-3 Zinc metalloproteinase-disintegrin-like berythracivase Zinc metalloproteinase-disintegrin-like bothrojarin-2 Zinc metalloproteinase-disintegrin-like brevilysin Zinc metalloproteinase-disintegrin-like cobrin Zinc metalloproteinase-disintegrin-like daborhagin-K Zinc metalloproteinase-disintegrin-like stejnihagin-A Zinc metalloproteinase-disintegrin-like stejnihagin-B Zinc metalloproteinase/disintegrin Zinc metalloproteinase/disintegrin Zinc metalloproteinase/disintegrin Metalloproteinase/disintegrin Zinc metalloproteinase/disintegrin Zinc metalloproteinase/disintegrin PMMP-1	Snake venom metalloproteinases
Cysteine-rich venom protein	Cysteine-rich venom protein
Disintegrin EO5A	Disintegrins

Table 4 (continued)

Proteins	Protein families
Disintegrin bitistatin	
L-amino-acid oxidase L-amino-acid oxidase L-amino-acid oxidase	L-amino acid oxidases
Long neurotoxin 1 Short neurotoxin 1 Weak toxin CM-2a Cytotoxin 1	Three-finger toxins
Snaclec 2 Snaclec 3 Snaclec CTL-Eoc124 Snaclec CTL-Eoc125 Snaclec agglucetin subunit beta-1 Snaclec bitiscetin subunit alpha Snaclec bitiscetin subunit beta Snaclec clone 2100755 Snaclec convulxin subunit beta C-type lectin 1	C-type lectins/Snaclecs
Snake venom 5'-nucleotidase	5' Nucleotidases
Snake venom vascular endothelial growth factor toxin barietin Vascular endothelial growth factor A	Vascular endothelial growth factors
Peroxiredoxin-4 Phospholipase A ₂ inhibitor 1 Snake venom glutaminyl-peptide cyclotransferase Hemoglobin subunit beta Secretory phospholipase A ₂ receptor Venom nerve growth factor 2	Peroxiredoxin Phospholipase A ₂ inhibitor Glutaminyl cyclases Globin Phospholipase A ₂ receptor Venom nerve growth factor

the ProteomeXchange Consortium via the PRIDE²⁶ partner repository with the dataset identifier PXD024638 and 10.6019/PXD024638, at <https://www.ebi.ac.uk/pride/archive/projects/PXD024638>.

Acknowledgements

Authors gratefully acknowledge Emeritus Professor Jose Maria Gutiérrez of the Clodomiro Picado Institute, University of Costa Rica, whose support, contributed tremendously to the success of this work. We would like to thank Professor Nicholas Casewell of the Liverpool School of Tropical Medicine for his valuable comments. Our appreciation goes to Dr. Peter Ofemile of the Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria, for his assistance in sourcing and milking of the snakes. This work did not receive any specific grants from public and private sectors.

References

- [1] J.M. Gutierrez, R.D. Theakston, D.A. Warrell, Confronting the neglected problem of 607 snakebite envenoming: the need for a global partnership, *PLoS Med.* 3 (2006) e150.
- [2] J.M. Gutiérrez, J.J. Calvete, A.G. Habib, R.A. Harrison, D.J. Williams, D.A. Warrell, Snakebite envenoming, *Nat. Rev. Dis. Primers* 3 (2017) 17063, <https://doi.org/10.1038/nrdp.2017.63>.
- [3] J.P. Chippaux, Snakebite envenomation turns again into a neglected tropical disease, *J. Venom. Anim. Toxins Incl. Trop. Dis.* 23 (2017) 38.
- [4] J.P. Chippaux, Estimate of the burden of snakebites in sub-Saharan Africa: a meta-analytic approach, *Toxicon* 57 (2011) 586–599.
- [5] D.A. Warrell, N.M.D. Davidson, B.M. Greenwood, L.D. Ormerod, H.M. Pope, B. J. Watkins, C.R.M. Prentice, Poisoning by bites of the saw-scaled or carpet viper (*Echis carinatus*) in Nigeria, *Q. J. Med.* 46 (1977) 33–62.
- [6] A.G. Habib, U.I. Gebi, G.C. Onyemelukwe, Snake bite in Nigeria, *Afr. J. Med. Med. Sci.* 30 (2001) 171–178.
- [7] A.G. Habib, Venomous snakes and snake envenomations in Nigeria, *Toxinol* 2 (2013) 275–298.
- [8] R. Ghosh, K. Mana, K. Gantait, S. Sarkhel, A retrospective study of clinico-epidemiological profile of snakebite related deaths at a tertiary care hospital in Midnapore, West Bengal, India, *Toxicol. Rep.* 5 (2018) 1–5.
- [9] B. Mohapatra, D.A. Warrell, W. Suraweera, P. Bhatia, N. Dhingra, R.M. Jotkar, P. S. Rodriguez, K. Mishra, R. Whitaker, P. Jha, Snakebite mortality in India: a

- nationally representative mortality survey, *PLoS Neglected Trop. Dis.* 5 (2011), e1018.
- [10] WHO, Snakebite envenoming: member States provide WHO with clear mandate for global action, Geneva, <https://www.who.int/news/item/25-05-2018-snakebite-envenoming-member-states-provide-who-with-clear-mandate-for-global-action>, 25th May, 2018. (Accessed November 2020).
- [11] S.B. Abubakar, I.S. Abubakar, A.G. Habib, A. Nasidi, N. Durfa, P.O. Yusuf, S. Larnyang, J. Garnvwa, E. Sokomba, L. Salako, G.D. Laing, R.D. Theakston, E. Juszczak, N. Alder, D.A. Warrell, Nigeria-UK EchiTab Study Group, Pre-clinical and preliminary dose-finding and safety studies to identify candidate antivenoms for treatment of envenoming by saw-scaled or carpet vipers (*Echis ocellatus*) in northern Nigeria, *Toxicon* 55 (2010) 719–723.
- [12] J.J. Calvete, Next-generation snake venomomics: protein-locus resolution through venom proteome decomplexation, *Expert Rev. Proteomics* 11 (2014) 315–329.
- [13] B. Lomonte, W.C. Tsai, J.M. Ureña-Díaz, L. Sanz, D. Mora-Obando, E.E. Sánchez, B. G. Fry, J.M. Gutiérrez, H.L. Gibbs, J.J. Calvete, Venomomics of New World pit vipers: genus-wide comparisons of venom proteomes across Agkistrodon, *J. Proteom.* 96 (2014) 103–116.
- [14] D. Pla, J.M. Gutiérrez, J.J. Calvete, Second generation snake antivenomics: comparing immunoaffinity and immunodepletion protocols, *Toxicon* 60 (2012) 688–699.
- [15] B.G. Fry, H. Scheib, L. Van DerWeerd, B. Young, J. McNaughtan, S.F. Ramjan, N. Vidal, R.E. Poelmann, J.A. Norman, Evolution of arsenal: structural and functional diversification of the venom system in advanced snakes (Caenophidia), *Mol. Cell. Proteomics* 7 (2008) 215–246.
- [16] T. Tasoulis, G.K. Isbister, A review and database of snake venom proteomes, *Toxins* 9 (2017) 673. E290.
- [17] J. Durban, L. Sanz, D. Trevisan-Silva, E. Neri-Castro, A. Alagon, J.J. Calvete, Integrated venomomics and venom gland transcriptome analysis of juvenile and adult Mexican rattlesnakes *Crotalus simus*, C. tzabcan, and C. culminatus revealed miRNA-modulated ontogenetic shifts, *J. Proteome Res.* 16 (2017) 3370–3390.
- [18] L. Goncalves-Machado, D. Pla, L. Sanz, R.J.B. Jorge, M. Leitao-De-Araujo, M. L. Alves, D.J. Alvares, J. De Miranda, J. Nowatzki, K. De Moraes-Zani, W. Fernandes, A.M. Tanaka-Azevedo, J. Fernandez, R.B. Zingali, J.M. Gutierrez, C. Correa-Netto, J.J. Calvete, Combined venomomics, venom gland transcriptomics, bioactivities, and antivenomics of two Bothrops jararaca populations from geographically isolated regions within the Brazilian Atlantic rainforest, *J. Proteomics* 135 (2016) 73–89.
- [19] A. Barlow, C.E. Pook, R.A. Harrison, W. Wuster, Coevolution of diet and prey-specific venom activity supports the role of selection in the evolution of snake venom, *Proc. Biol. Sci.* 276 (2009) 2443–2449.
- [20] G.S. Dias, E.S. Kitano, A.H. Pagotto, S.S. Sant'anna, M.M. Rocha, A. Zelanis, S. M. Serrano, Individual variability in the venom proteome of juvenile *Bothrops jararaca* specimens, *J. Proteome Res.* 12 (2013) 4585–4598.
- [21] S. Kunalan, I. Othman, S.S. Hassan, W.C. Hodgson, Proteomic characterization of two medically important Malaysian snake venoms, *Calloselasma rhodostoma* (Malayan pit viper) and *Ophiophagus hannah* (king Cobra), *Toxins* 10 (2018), <https://doi.org/10.3390/toxins10110434>.
- [22] N.R. Casewell, S.C. Wagstaff, W. Wüster, D.A.N. Cook, F.M.S. Bolton, S.I. King, D. Pla, L. Sanz, J.J. Calvete, R.A. Harrison, Postgenomic processes dictate venom variation, *Proc. Natl. Acad. Sci. Unit. States Am.* 111 (2014) 9205–9210.
- [23] S.C. Wagstaff, R.A. Harrison, Venom gland EST analysis of the saw-scaled viper, *Echis ocellatus*, reveals novel alpha9beta1 integrin-binding motifs in venom metalloproteinases and a new group of putative toxins, renin-like aspartic proteases, *Gene* 377 (2006) 21–32.
- [24] R.E. Hill, S.P. Mackessy, Venom yields from several species of colubrid snakes and differential effects of ketamine, *Toxicol.* 35 (1997) 671–678.
- [25] F.A. Adamude, E.J. Dingwoke, M.S. Abubakar, S. Ibrahim, G. Mohamed, A. Klein, A.B. Sallau, Proteomic analysis of three medically important Nigerian *Naja* (*Naja haje*, *Naja katiensis* and *Naja nigricollis*) snake venoms, *Toxicol.* 197 (2021) 24–32.
- [26] B. Lomonte, W.C. Tsai, J.M. Ureña-Díaz, L. Sanz, D. Mora-Obando, E.E. Sánchez, B. G. Fry, J.M. Gutiérrez, H.L. Gibbs, M.G. Sovic, J.J. Calvete, Venomomics of New World pit vipers: genus-wide comparisons of venom proteomes across Agkistrodon, *J. Proteomics* 96 (2014) 103–116.
- [27] H. Liu, R.G. Sadygov, J.R. Yates, A model for random sampling and estimation of relative protein abundance in shotgun proteomics, *Anal. Chem.* 76 (2004) 4193–4201.
- [28] Y. Zhang, Z. Wen, M.P. Washburn, L. Florens, Refinements to label free proteome quantitation: how to deal with peptides shared by multiple proteins, *Anal. Chem.* 82 (2010) 2272–2281.
- [29] D. Fermin, V. Basrur, A.K. Yocum, A.I. Nesvizhskii, Abacus: a computational tool for extracting and pre-processing spectral count data for label-free quantitative proteomic analysis, *Proteomics* 11 (2011) 1340–1345.
- [30] Y. Perez-Riverol, A. Csordas, J. Bai, M. Bernal-Llinares, S. Hewapathirana, D.J. Kundu, A. Inuganti, J. Griss, G. Mayer, M. Eisenacher, E. Perez, J. Uszkoreit, J. Pfeuffer, T. Sachsenberg, S. Yilmaz, S. Tiwary, J. Cox, E. Audain, M. Walzer, A.F. Jarnuczak, T. Ternent, A. Brazma, J.A. Vizcaino, The PRIDE database and related tools and resources in 2019: improving support for quantification data, *Nucleic Acids Res.* 47 (D1): D442–D450.
- [31] D. H. Lundgren, S. Hwang, L. Wu, D.K. Han, Role of spectral counting in quantitative proteomics, *Expert Rev. Proteomics* 7 (2010) 39–53.
- [32] M. Blein-Nicolas, M. Zivy, Thousand and One Ways to Quantify and Compare Protein Abundances in Label-free Bottom-Up Proteomics, *Biochim Biophys Acta Proteom. Proteom.* 2016, <https://doi.org/10.1016/j.bbapap.2016.02.019>.
- [33] J.M. Gutierrez, A. Rucavado, T. Escalante, C. Dias, Hemorrhage induced by snake venom metalloproteinases: biochemical and biophysical mechanisms involved in microvessel damage, *Toxicon* 45 (2005) 997–1011.
- [34] J.J. Calvete, P. Juarez, L. Sanz, Snake venomomics. Strategy and applications, *J. Mass Spectrom.* 42 (2007) 1405–1414.
- [35] N.R. Casewell, R.A. Harrison, W. Wüster, S.C. Wagstaff, Comparative venom gland transcriptome surveys of the saw-scaled vipers (*Viperidae: Echis*) reveal substantial intra-family gene diversity and novel venom transcripts, *BMC Genom.* 10 (2009) 564, <https://doi.org/10.1186/1471-2164-10-564>.
- [36] T.S. Kang, D. Georgieva, N. Genov, M.T. Murakami, M. Sinha, R.P. Kumar, P. Kaur, W.S. Kurmar, S. Dey, S. Sharma, A. Vrieling, C. Betzel, S. Takeda, R.K. Arni, T. P. Singh, R.M. Kini, Enzymatic toxins from snake venoms: structural characterization and mechanism of catalysis, *FEBS J.* 278 (2011) 4544–4576.
- [37] T. Sajevic, A. Leonardi, I. Krizaj, Haemostatically active proteins in snake venoms, *Toxicon* 57 (2011) 627–645.
- [38] S. Waidyanatha, A. Silva, S. Siribaddana, G.K. Isbister, Long-term effects of snake envenoming, *Toxins* 11 (2019) 193, <https://doi.org/10.3390/toxins11040193>.
- [39] D.A. Warrell, Snake bite, *Lancet* 375 (2010) 77–88.
- [40] J.M. Gutierrez, A. Rucavado, T. Escalante, C. Dias, Hemorrhage induced by snake venom metalloproteinases: biochemical and biophysical mechanisms involved in microvessel damage, *Toxicon* 45 (2005) 997–1011.
- [41] A.M. Moura-da-Silva, M.T. Almeida, J.A. Portes-Junior, C.A. Nicolau, F. Gomes-Neto, R.H. Valente, Processing of snake venom metalloproteinases: generation of toxin diversity and enzyme inactivation, *Toxins* 8 (2016) 183, <https://doi.org/10.3390/toxins8060183>.
- [42] M. Kohlhoff, M.H. Borges, A. Yarleque, C. Cabezas, M. Richardson, E.F. Sanchez, Exploring the proteomes of the venoms of the Peruvian pit vipers, *J. Proteom.* 75 (2012) 2181–2195.
- [43] J. White, Snake venom and coagulopathy, *Toxicol.* 45 (2005) 951–967.
- [44] N.R. Casewell, W. Wüster, F.J. Vonk, R.A. Harrison, B.G. Fry, Complex cocktails: the evolutionary novelty of venoms, *Trends Ecol. Evol.* 28 (2013) 219–229.
- [45] J.J. Calvete, Venomomics: integrative venom proteomics and beyond, *Biochem. J.* 474 (2017) 611–634.
- [46] N.R. Casewell, On the ancestral recruitment of metalloproteinases into the venom of snakes, *Toxicol.* 60 (2012) 449–454.
- [47] F.S. Markland-Jnr, S. Swenson, Snake venom metalloproteinases, *Toxicol.* 62 (2013) 3–18.
- [48] T. Escalante, A. Rucavado, J.W. Fox, J.M. Gutierrez, Key events in microvascular damage induced by snake venom hemorrhagic metalloproteinases, *J. Proteom.* 74 (2011) 1781–1794.
- [49] S. Takeda, H. Takeya, S. Iwanaga, Snake venom metalloproteinases: structure, function and relevance to the mammalian ADAM/ADAMTS family proteins, *Biochim. Biophys. Acta* 1824 (2012) 164–176.
- [50] J. Slagboom, J. Kool, R.A. Harrison, N.R. Casewell, Haemotoxic snake venoms: their functional activity, impact on snakebite victims and pharmaceutical promise, *Br. J. Haematol.* 177 (2017) 947–959.
- [51] C.C. Hsu, W.B. WB, T.F. Huang, A snake venom metalloproteinase, kistomin, cleaves platelet glycoprotein VI and impairs platelet functions, *J. Thromb. Haemostasis* 6 (2008) 1578–1585.
- [52] E.K. Johnson, C.L. Ownby, Isolation of a hemorrhagic toxin from the venom of *Agkistrodon contortrix laticinctus* (Broad-banded copperhead) and pathogenesis of the hemorrhage induced by the toxin in mice, *Int. J. Biochem.* 25 (1993) 267–278.
- [53] W.J. Wang, T.F. Huang, Purification and characterization of a novel metalloproteinase, acurhagin, from *Agkistrodon acutus* venom, *Thromb. Haemostasis* 87 (2002) 641–650.
- [54] T. Nikai, K. Taniguchi, Y. Komori, K. Masuda, J.W. Fox, H. Sugihara, Primary structure and functional characterization of biltoxin-1, a novel dimeric P-II snake venom metalloproteinase from *Agkistrodon bilineatus* venom, *Arch. Biochem. Biophys.* 378 (2000) 6–15.
- [55] E.F. Sanchez, L.M. Gabriel, S. Gontijo, L.H. Gremiski, S.S. Veiga, K.S. Evangelista, J. A. Eble, M. Richardson, Structural and functional characterization of a P-III metalloproteinase, leucurolysin-B, from *Bothrops leucurus* venom, *Arch. Biochem. Biophys.* 468 (2007) 193–204.
- [56] L.G. Jia, X.M. Wang, J.D. Shannon, J.B. Bjarnason, J.W. Fox, Function of disintegrin-like/cysteine-rich domains of Atrolysin A, *J. Biol. Chem.* 272 (1997) 13094–13102.
- [57] M.B. Silva, M. Schattner M, C.R. Ramos, I.L. Junqueira-de-Azevedo, M. C. Guarnieri, M.A. Lazzari, C.A. Sampaio, R.G. Pozner, J.S. Ventura, P. L Ho, A. M. Chudzinski-Tavassi, A prothrombin activator from *Bothrops erythromelas* (jararaca-da-seca) snake venom: characterization and molecular cloning, *Biochem. J.* 369 (2003) 129–139.
- [58] S.M.T. Serrano, The long road of research on snake venom serine proteinases, *Toxicol.* 62 (2013) 19–26.
- [59] K. Bharati, S.S. Hasson, J. Oliver, G.D. Laing, R.D.G. Theakston, R.A. Harrison, Molecular cloning of phospholipases A₂ from venom glands of *Echis* carpet vipers, *Toxicol.* 41 (2003) 941–947.
- [60] E. Jiménez-Charris, L. Montealegre-Sanchez, L. Solano-Redondo, D. Mora-Obando, E. Camacho, F. Castro-Herrera, L. Fierro-Pérez, B. Lomonte, Proteomic and functional analyses of the venom of *Porthidium lansbergii lansbergii* (Lansberg's hognoe viper) from the Atlantic Department of Colombia, *J. Proteom.* 114 (2015) 287–299.
- [61] J.B. Harris, B.D. Grubb, C.A. Maltin, R. Dixon, The neurotoxicity of the venom phospholipases A₂, notexin and taipoxin, *Exp. Neurol.* 161 (2000) 517–526.
- [62] J.B. Harris, T. Scott-Davey, Secreted phospholipases A₂ of snake venoms: effects on the peripheral neuromuscular system with comments on the role of phospholipases A₂ in disorders of the CNS and their uses in industry, *Toxins* 5 (2013) 2533–2571.
- [63] J.M. Gutierrez, C.L. Ownby, Skeletal muscle degeneration induced by venom phospholipases A₂: insights into the mechanisms of local and systemic myotoxicity, *Toxicol.* 42 (2003) 915–931.

- [64] Y. Angulo, J. Escolano, B. Lomonte, J.M. Gutiérrez, L. Sanz, J.J. Calvete, Snake venomomics of central American Pitvipers: Clues for rationalizing the distinct envenomation profiles of *Atropoides nummifer* and *Atropoides picadoi*, *J. Proteome Res.* 7 (2008) 708–719.
- [65] N.A. Ghazaryan, L. Ghulikyan, A. Kishmiryan, T.V. Andreeva, Y.N. Utkin, V. I. Tsetlin, B. Lomonte, N.M. Ayvazyan, Phospholipases a2 from Viperidae snakes: differences in membranotropic activity between enzymatically active toxin and its inactive isoforms, *Biochim. Biophys. Acta Biomembr.* 1848 (2015) 463–468.
- [66] K. Trummal, M. Samel, A. Aaspõllu, K. Tõnismägi, T. Titma, J. Subbi, J. Siigur, E. Siigur, 5'-Nucleotidase from *Vipera lebetina* venom, *Toxicon* 93 (2015) 155–163.
- [67] S.D. Aird, Ophidian envenomation strategies and the role of purines, *Toxicon* 40 (2002) 335–393.
- [68] B.L. Dhananjaya, C.J. D'Souza, The pharmacological role of nucleotidases in snake venoms, *Cell Biochem. Funct.* 28 (2010) 171–177.
- [69] X.Y. Du, K.J. Clemetson, Snake venom L-amino acid oxidases, *Toxicon* 40 (2002) 659–665.
- [70] S. Torii, M. Naito, T. Tsuruo, I. Apoxin, A novel apoptosis-inducing factor with L-amino acid oxidase activity purified from Western diamondback rattlesnake venom, *J. Biol. Chem.* 272 (1997) 9539–9542.
- [71] J.W. Fox, S.M. Serrano, Insights into and speculations about snake venom metalloproteinase (SVMP) synthesis, folding and disulfide bond formation and their contribution to venom complexity, *FEBS J.* 275 (2008) 3016–3030.
- [72] Y. Komori, T. Nikai, T. Tohkai, H. Sugihara, Primary structure and biological activity of snake venom lectin (APL) from *Agkistrodon p. piscivorus* (Eastern cottonmouth), *Toxicon* 37 (1999) 1053–1064.
- [73] Y. Ozeki, T. Matsui, J. Hamako, M. Suzuki, Y. Fujimura, E. Yoshida, S. Nishida, K. Titani, C-type galactoside-binding lectin from *Bothrops jararaca* venom: comparison of its structure and function with those of botrocetin, *Arch. Biochem. Biophys.* 308 (1994) 306–310.
- [74] P. Kessler, P. Marchot, M. Silva, D. Servent, The three-finger toxin fold: a multifunctional structural scaffold able to modulate cholinergic functions, *J. Neurochem.* 142 (2017) 7–18.
- [75] C.M. Barber, G.K. Isbister, W.C. Hodgson, Alpha neurotoxins, *Toxicon* 66 (2013) 47–58.
- [76] D. Chakrabarty, C. Chanda, Snake venom disintegrins, in: P. Gopalakrishnakone, H. Inagaki, C.W. Vogel, A. Mukherjee, T. Rahmy (Eds.), *Snake Venoms*, *Toxinol*, Springer, Dordrecht, 2017, https://doi.org/10.1007/978-94-007-6410-1_14.
- [77] S. Niewiarowski, M.A. MacLane, M. Klocsewiak, G.J. Stewart, Disintegrins and other naturally occurring antagonists of platelet fibrinogen receptor, *Semin. Hematol.* 31 (1994) 289–300.
- [78] T. Kostiza, J. Meier, Nerve growth factors from snake venoms: Chemical properties, mode of action and biological significance, *Toxicon* 34 (1996) 787–806.
- [79] R.H. Ziganshin, S.I. Kovalchuk, G.P. Arapidi, V.G. Starkov, A.N. Hoang, T.T. T. Nguyen, K.C. Nguyen, B.B. Shoibonov, V.I. Tsetlin, Y.N. Utkin, Quantitative proteomic analysis of Vietnamese krait venoms: neurotoxins are the major components in *Bungarus multicinctus* and phospholipases A2 in *Bungarus fasciatus*, *Toxicon* 107 (2015) 197–209.
- [80] S.I. Kovalchuk, R.H. Ziganshin, V.G. Starkov, V.I. Tsetlin, Y.N. Utkin, Quantitative proteomic analysis of venoms from Russian vipers of *Pelias* group: phospholipases A2 are the main venom components, *Toxins* 8 (2016) 105, <https://doi.org/10.3390/toxins8040105>.
- [81] H. Xiao, H. Pan, K. Liao, M. Yang, C. Huang, Snake venom PLA2, a promising target for broad-spectrum antivenom drug development, *BioMed Res. Int.* (2017), <https://doi.org/10.1155/2017/6592820>.
- [82] N.A. Santos-Filho, C.T. Santos, Alpha-type phospholipase A2 inhibitors from snake blood, *J. Venom. Anim. Toxins Incl. Trop. Dis.* (2017), <https://doi.org/10.1186/s40409-017-0110-2>.
- [83] P.C. Campos, L.A. de Melo, G.L.F. Dias, C.L. Fortes-Dias, Endogenous phospholipase A₂ inhibitors in snakes: a brief overview, *J. Venom. Anim. Toxins Incl. Trop. Dis.* 21 (2016) 37, 22.